arXiv:1207.2383v2 [astro-ph.GA] 27 Jul 2012

The Temperature of Dark Matter Atmospheres

Sanjoy K. Sarker* and Allen Stern†

Department of Physics, University of Alabama,
Tuscaloosa, Alabama 35487, USA

ABSTRACT

We model the dark matter in galactic haloes with a self-gravitating atmosphere surrounding the galaxy. The galaxy serves to set the scales and the boundary conditions for the atmosphere. The atmosphere is treated as an isothermal Boltzmann gas, which at sufficiently large distances leads to flat rotation curves. Solutions to the dynamics are determined by two parameters, one of which is the ratio of the dark matter mass to the equilibrium temperature. From typical orbital speeds in haloes, any dark matter candidate that utilizes this mechanism to generate flat rotation curves should have a mass to temperature ratio of ~ 400 eV per degree Kelvin.

---

* ssarker@bama.ua.edu
†astern@bama.ua.edu

The system of a galaxy plus halo is often approximated at zeroth order exclusively in terms of its dominant dark matter component. A simple model, which has been around for quite some time, assumes that all the dark matter particles are in thermal equilibrium and form a self-gravitating Boltzmann gas.[1],[2] The dynamics of this system is governed by the Emden equation.[3] The standard spherically symmetric solution is obtained by imposing that the gravitational force vanishes at the origin $r = 0$. The resulting dark matter density profile has a $1/r^2$ behavior beyond a certain distance scale known as the King radius[1], which is generally associated with the galaxy core. This is just what is required for objects undergoing circular orbits in the medium to have an r-independent orbital speed. (Currently many more density profiles are available, such as the NFW profile[5], which are capable of producing approximately flat rotation curves at some distance scales. The profiles have been fit to observations in galactic haloes, and also to gravitational lensing effects.[6],[7],[8]) The above approach assumes that the Boltzmann gas approximation is valid for all r, including distances that are small compared with the King radius; i.e. corresponding to being within the galaxy. The more involved thermodynamic behavior of the baryonic component to the galaxy, which interacts gravitationally with the gas, is ignored in this zeroth order treatment. Thus baryonic matter is to be treated as a perturbation in this approach. Complications can then arise in the short distance regime, as baryonic matter plays an important role there, and this in turn can effect the large distance behavior of the gas. Complications at short distances also plague dark matter simulations, where cusps appear in the density profile at $r = 0$.[‡][5] Such singularities do not agree with observations. (In [9] it was shown that the cusps may be eliminated by taking into account the quantum statistics of fermions.)

A more realistic picture should result if one takes into account the baryonic component in some manner from the outset. Its presence introduces a new distance scale, which we denote by R_G, as well as a mass scale M_G, which is the mass of the galaxy. R_G defines a boundary between the galactic inner region (with a low mass to luminosity ratio) and the halo outer region (with a high mass to luminosity ratio). We refer to the latter as the galactic atmosphere. (We shall discuss the relation of R_G to the King radius later.) Here we shall not be concerned with the details of the system for $r < R_G$. In our model, the interior region only serves to fix the boundary conditions for the atmosphere, for which a simple Boltzmann gas description now appears more reasonable. This is because densities become low enough for the system to behave as a classical gas at sufficiently large distances from the origin. This gives an operational definition of the distance scale R_G, i.e., it is the distance scale beyond which the isothermal gas approximation is valid. The Emden equations can then be employed to describe the gas in the region. Here the boundary conditions should be imposed at $r = R_G$, rather than at the origin. The boundary conditions allow for a family of solutions to the Emden equations, which are parametrized by constants depending on M_G and R_G. The family of solutions contain the standard spherically symmetric solution discussed above, i.e., the one obtained by demanding that the gravitational force vanishes at the origin, as a special case. The resulting density profiles should then provide a better fit with observations than that associated with the standard spherically symmetric solution. As before, flat rotation curves result beyond a certain distance, which we denote by $\mathcal{R} \geq R_G$. Among the solutions is a well known one which gives a flat rotation curve for the entire domain. The solution has been referred to as the singular isothermal sphere because it is singular at $r = 0$ if one assumes that the isothermal Boltzmann gas approximation is valid for all r. As stated above, such an assumption is suspect in view of the complicated dynamics within the galaxy, and we shall not presume that it holds. Here instead the solution is *nonsingular* because the domain is restricted to the galactic atmosphere, $r \geq R_G$. It results from particular values of the two parameters, one of which is the ratio of the dark particle mass to temperature, m/T. The associated flat rotation curve for all $r \geq R_G$ means that this solution serves as a crude fit to observations. From this fit we can obtain an estimate of the two parameters. Upon substituting typical orbital speeds of ~ 200 km/s found in the haloes, we find $m/T \sim 400$ eV per degree Kelvin of the atmosphere. This result does not allow for hot dark matter with particle mass below ~ 800 eV, since the $\sim 2\ ^o$K relic background temperature serves as a lower bound for the equilibrium temperature of the halo. Aside from this remark, it is not our intention to favor any particular dark matter candidate, but rather to examine universal features resulting from this simple model of galactic haloes.

[‡] Temperature has been introduced in numerical simulations, and it was claimed to be an $r-$dependent function.[4]

In what follows we give a description of the model and some simple implications of the results. We begin with the basic thermodynamics of a self-gravitating isothermal gas. It results from the balancing of the kinetic pressure of the gas with the gravitational attraction to the galaxy and to itself. While thermal equilibrium may not be a good approximation within a galaxy, it most likely can be trusted to give a reasonable picture of the large-scale structure beyond the galaxy, and so we consider the restricted domain $r \geq R_G$. We shall neglect nonequilibrium processes that may occur near $r = R_G$ (and also beyond some much larger scale where an atmosphere may no longer be viable). Here we shall assume a nonrelativistic ideal gas[§] consisting of only one type of particle[¶], with mass m, moving in a slowly varying[‖] gravitational potential ϕ. The geometry of the galaxy should become less important as the distance from it increases. Therefore, it is reasonable to assume that spherical symmetry holds at lowest order, and so quantities of interest, such as ϕ and the average number density in the atmosphere n_{atm}, are functions of only the radial variable. In general, the average number density has the form $n_{atm}(r) = n(\mu - m\phi(r))$, μ being the chemical potential. For a Boltzmann gas, $n(\mu) = e^{\frac{\mu}{k_B T}}/\Lambda^3$, where $\Lambda = \sqrt{\frac{2\pi\hbar^2}{mk_B T}}$ is the thermal wavelength and T is the temperature. The value of ϕ at $r = R_G$ is arbitrary. It is convenient to choose it to be zero so that the number density is $n(\mu)$ at $r = R_G$, and

$$n_{atm}(r) = n_{atm}(R_G) e^{-m\phi(r)/k_B T} \ . \tag{1}$$

On the other hand, the value of $\frac{d\phi}{dr}$ is fixed at $r = R_G$ since it is the acceleration of gravity at $r = R_G$. Thus

$$\phi(R_G) = 0 \qquad\qquad \frac{d\phi}{dr}\Big|_{r=R_G} = \frac{GM_G}{R_G^2} \ , \tag{2}$$

M_G being the galactic mass. From the two boundary conditions we can then solve the Poisson equation for $\phi(r)$,

$$\frac{1}{r^2}\frac{d}{dr}\Big(r^2\frac{d\phi}{dr}\Big) = 4\pi Gm\, n_{atm}(R_G) e^{-m\phi(r)/k_B T} \ , \qquad\qquad r \geq R_G \tag{3}$$

After a rescaling to the dimensionless variables $x = \frac{r}{R_G}$ and $\chi = \frac{R_G}{GM_G}\phi$, the equation simplifies to

$$\frac{1}{x^2}\frac{d}{dx}\Big(x^2\frac{d\chi}{dx}\Big) = 3\kappa\, e^{-\frac{\chi(x)}{\tau}} \ , \qquad\qquad x \geq 1 \tag{4}$$

along with boundary conditions

$$\chi(1) = 0 \qquad\qquad \frac{d\chi}{dx}\Big|_{x=1} = 1 \tag{5}$$

[§] We shall verify that the relativistic corrections are small. Also, we shall find the appropriate range for the dark particle mass where the approximation of a Boltzmann gas is valid and the effects of quantum statistics can be ignored.

[¶] In an atmosphere consisting of several components (including dark matter particles) in equilibrium, the heavier particles will sink toward the boundary region at $r = R_G$, while the long range behavior is governed by the lightest particles.

[‖] i.e., slowly varying with respect to the average interparticle distance

Here κ and τ are independent dimensionless parameters determining the solutions. They depend, among other things, on R_G and M_G. κ is the ratio of the mean mass density of the atmosphere at $r = R_G$ to the mean mass density of the galaxy $\rho_G = \frac{M_G}{\frac{4}{3}\pi R_G^3}$, while τ involves m, T and the speed $v_{x=1} = \sqrt{GM_G/R_G}$ of objects undergoing circular orbits at $r = R_G$,

$$\kappa = \frac{mn(\mu)}{\rho_G} \qquad\qquad \tau = \frac{k_B T}{m v_{x=1}^2} \tag{6}$$

Quantities of interest are the speed v of objects undergoing circular orbits in the atmosphere as a function of the radial coordinate and the mass M_{atm} of the atmosphere enclosed in a sphere of radius $r > R_G$. They are given respectively by

$$\left(\frac{v}{v_{x=1}}\right)^2 = x\frac{d\chi}{dx} \qquad \text{and} \qquad \frac{M_{atm}(x)}{M_G} = 3\kappa\int_1^x dx' x'^2 e^{-\frac{\chi(x')}{\tau}} \tag{7}$$

Eq. (4) is the spherically symmetric Lane-Emden equation for infinite polytropic index, which is often referred to as the Emden equation.[3] Various techniques have been employed to obtain their solutions, and their application to galactic dynamics has been discussed. (See for example, [2],[10],[11],[12]. Axially symmetric solutions with rotation have also been obtained.[13]) The boundary conditions for this equation are standardly imposed at the origin, where the gravitational force is required to vanish $\frac{d\chi}{dx}\Big|_{x=0} = 0$.** As mentioned above, the validity of the solutions for $x < 1$ is not so clear in application to galaxies. This is not a concern here, since our domain is $x \geq 1$. Having the boundary conditions specified at $x = 1$, i.e., at the boundary of the galaxy and atmosphere, allows us to avoid complications in the interior region. In this model, the interior region only serves to generate the external gravitational field in the atmosphere.

Solutions to the Emden equations generated from boundary conditions (5) at $x = 1$ form a larger family than those resulting from the boundary conditions at the origin stating that the gravitational force vanishes there. More specifically, solutions that can be consistently continued from $x = 1$ to the origin, where $\frac{d\chi}{dx}\Big|_{x=0} = 0$, form a subset of the family of solutions satisfying (5). κ and τ are not independent for the subset. To see this one can do another rescaling of the field and the coordinate, $\chi \to \tilde{\chi} = \chi/\tau$ and $x \to \tilde{x} = \sqrt{\frac{3\kappa}{\tau}}\, x$, removing the parameters from the differential equation (4), and putting them instead in the boundary conditions. Then

$$\frac{1}{\tilde{x}^2}\frac{d}{d\tilde{x}}\left(\tilde{x}^2\frac{d\tilde{\chi}}{d\tilde{x}}\right) = e^{-\tilde{\chi}(\tilde{x})}\ , \tag{8}$$

with

$$\tilde{\chi}\Big|_{\tilde{x}=\sqrt{\frac{3\kappa}{\tau}}} = 0 \qquad\qquad \frac{d\tilde{\chi}}{d\tilde{x}}\Big|_{\tilde{x}=\sqrt{\frac{3\kappa}{\tau}}} = \frac{1}{\sqrt{3\kappa\tau}} \tag{9}$$

For generic values of κ and τ, the boundary conditions (9) cannot be generated by integrating the Emden equation from the origin, starting from $\frac{d\tilde{\chi}}{d\tilde{x}}\Big|_{\tilde{x}=0} = 0$ and any $\tilde{\chi}(0)$. [$\tilde{\chi}(0)$ must be negative since $\tilde{\chi}$ grows with $\tilde{x}$.] Equivalently, if one were to impose the boundary condition $\frac{d\tilde{\chi}}{d\tilde{x}}\Big|_{\tilde{x}=0} = 0$, along with some $\tilde{\chi}(0) < 0$, one could integrate the Emden equations to find the intersection of $\tilde{\chi}(\tilde{x})$ with the $\tilde{x}-$axis, along with the slope of the function at that point. One can then use (9) to numerically solve for specific values of κ and τ, defining the subset of solutions. The King radius[1], which is the radius where the dark matter density is half the density at $r = 0$, can be defined for these special values of κ

** χ is often set to zero, as well, at the origin, but this is a gauge condition which can be removed with a redefinition of κ.

and τ. It is $r_{\rm King} = \frac{R_G}{2}\sqrt{\frac{3\tau}{\kappa}}\sqrt{\frac{n(\mu)}{n_{atm}(0)}}$, where $n_{atm}(0)$ is the number density of the gas continued to the origin. From (1) it is $n(\mu)e^{-\tilde{\chi}(0)}$, and so

$$r_{\rm King} = \frac{R_G}{2}\sqrt{\frac{3\tau}{\kappa}}\, e^{\tilde{\chi}(0)/2} \tag{10}$$

This expression is only valid for special values of κ and τ, determined from $\tilde{\chi}(0)$ using (9). As a result of the latter, $r_{\rm King}/R_G$ is a function only of $\tilde{\chi}(0)$. In contrast, our approach makes no apriori restrictions on the values of κ and τ, and the behavior of the system for $r < R_G$ is not assumed to be governed by the Emden equations. The King radius cannot be defined for generic values of κ and τ.

Only one analytic solution to the Emden equations (4) is known and it is[1]

$$\chi(x) = \tau \log\left(\frac{3\kappa x^2}{2\tau}\right) \tag{11}$$

It is called the 'singular' isothermal sphere because it is ill-defined at the origin (for any values of κ and τ). Rather than satisfying $\frac{d\chi}{dx}\Big|_{x=0} = 0$, the acceleration of gravity diverges at the origin. On the other hand, (11) is mathematically and physically well defined for the domain of interest here, $x \geq 1$. It satisfies the boundary conditions (5) for the case of $\tau = \frac{1}{2}$ and $\kappa = \frac{1}{3}$, or equivalently,

$$n_{atm}(R_G) = \frac{M_G}{4\pi m R_G^3} \qquad\qquad \frac{m}{T} = \frac{2k_B}{v_{x=1}^2} \tag{12}$$

In this case, (7) states that the orbital speed is independent of x, $v = v_{x=1}$. It is interesting to note that this orbital speed is of the same order of magnitude as the rms speed u of the constituent particles of the gas. For a monatomic gas, $u = \sqrt{\frac{3}{2}}v$. From (7) one also gets a linear function for M_{atm}: $M_{atm}(x)/M_G = x - 1$.

Eq. (11) is obviously not a solution for arbitrary values of τ and κ. However, (11) does describe the large distance behavior of generic solutions††, and satisfies the physically desired feature that the gravitational force tends to zero as $x \to \infty$. (The potential is singular in this limit.) The orbital speed goes to the constant value, $v = \sqrt{2\tau}\, v_{x=1}$ (or zero, as in the case of no atmosphere $\kappa = 0$) for large x. This is illustrated for various values of τ and κ [including the values $\tau = \frac{1}{2}$ and $\kappa = \frac{1}{3}$ associated with the exact solution (11)] in figures 1 and 2. $M_{atm}(x)/M_G$ tends to a linear function of x with slope approaching 2τ as $x \to \infty$, which is illustrated in figures 3 and 4. The asymptotic region $x \geq \mathcal{R}/R_G \geq 1$, where solutions tends to (11), depends on the values of κ and τ. The asymptotic region becomes the entire domain, $x \geq 1$, in the limit that (11) is exact, $\tau \to \frac{1}{2}$ and $\kappa \to \frac{1}{3}$. Since approximately flat rotation curves are observed in galactic haloes for all measured values of the radial coordinate beyond a certain distance scale (which we can identify with R_G), then solution (11) is approximately valid for all $x > 1$. So $\tau \approx \frac{1}{2}$ and $\kappa \approx \frac{1}{3}$ roughly holds for all galaxies and (12) should provide a crude estimate for $n(\mu)$ and m/T. From the former, the mass density of the galactic atmosphere at the boundary $x = 1$ is approximately one-third the mass density of the galaxy.

We can now check the validity of the nonrelativistic Boltzmann approximation which was used here. As we found earlier, the rms speed u of the constituent particles of the gas is of the same order of magnitude as v. A typical value of ~ 200 km/s can be used for the r-independent speed $v = v_{x=1}$ in haloes, which serves as an order of magnitude estimate for u.[2] It follows that relativistic corrections for the gas are small, although not completely negligible. (In this sense, any dark matter particle making

---

†† For this we can do another change of coordinates and rewrite the differential equation in terms of $y = \log x$. Upon assuming that $\frac{d^2\chi}{dy^2} << \frac{d\chi}{dy}$ when $y \to \infty$, one gets $\frac{d\chi}{dy} \sim 3\kappa \exp\left(2y - \chi(y)/\tau\right)$, the solution of which is (11).



Figure 1: $v/v_{x=1}$ is plotted versus x for $\tau = \frac{1}{2}$ and $\kappa = 0, \frac{1}{6}, \frac{1}{3}$ and $\frac{1}{2}$. The values of κ are listed in ascending order of the slope at $x = 1$, with $\kappa = 0$ corresponding to no atmosphere.



Figure 2: $v/v_{x=1}$ is plotted versus x for $\kappa = \frac{1}{3}$ and $\tau = \frac{1}{4}, \frac{1}{2}$ and 1. The values of τ are listed in ascending order of the slope at $x = 1$. The limiting value of $v/v_{x=1}$ is $\sqrt{2\tau}$.

up such a galactic atmosphere can be considered *warm.*) Our approximation of a classical Boltzmann gas is valid provided that the thermal wavelength is much smaller than the interparticle distance, or

$$\frac{mk_BT}{2\pi\hbar^2 n_{atm}^{2/3}} >> 1 \ , \tag{13}$$

with the smallest value of the left-hand-side occurring at the boundary $r = R_G$. Upon using (12), this requires that $m^{8/3} >> 4\pi\hbar^2(\rho_G/3)^{2/3}/v_{x=1}^2$. For an order of magnitude estimate, we take $\rho_G \sim 10^9 M_\odot/\text{Kpc}^3$ along with $v_{x=1} \sim 200$ km/s, to get the result that the Boltzmann approximation is valid provided that dark particle mass is significantly greater than ~ 100 eV. Quantum statistics is therefore relevant for galactic atmospheres consisting of dark matter particles with $m < 100$ eV, however such hot dark matter scenarios are currently disfavored. (Similar conclusions appear in [5].)

Upon substituting $v_{x=1} \sim 200$ km/s in (12) one ends up with an estimate for m/T of about 400 eV per Kelvin. Since the ~ 2 ^{o}K background temperature provides a lower limit on the equilibrium temperature for any dark matter relic, the particle constituent making up the isothermal atmosphere cannot have a mass far below 800 eV, which is consistent with the case against hot dark matter particles. (A similar argument was used to estimate a lower bound on the sterile neutrino mass in [2].) The value



Figure 3: M_{atm}/M_G is plotted versus x for $\tau = \frac{1}{2}$ and $\kappa = \frac{1}{6}, \frac{1}{3}$ and $\frac{1}{2}$. The values of κ are listed in ascending order of the slope at $x = 1$.



Figure 4: M_{atm}/M_G is plotted versus x for $\kappa = \frac{1}{3}$ and $\tau = \frac{1}{4}, \frac{1}{2}$ and 1. The values of τ are listed in ascending order of the slope at $x = 1$. The limiting value of the slope is 2τ as $x \to \infty$.

of 800 eV is generally considered below the current lower bound for warm dark matter mass (see for example, [14],[15]), and so galactic atmospheres should have an equilibrium temperature above the $\sim 2\ {}^o$K background.

The model discussed here can be applied to any dark matter candidate with mass greater than $\sim$ 800 eV. A promising candidate for warm dark matter is the sterile neutrino whose mass is thought to be in the KeV range. (For a small sample of articles, see [16],[17],[18],[19],[20],[21],[22]. The possibility of neutral leptons being dark matter candidates was considered much earlier in [23].) For the case of galactic atmospheres consisting of 2 KeV particles, the associated equilibrium temperature should be $\sim 5\ {}^o$K. At the other end of the spectrum, an atmosphere consisting of cold dark matter particles would require an enormous temperature to sustain the desired density profile. For the example of a 100 GeV particle one gets temperatures of the order $10^{8-9}\ {}^o$K. At at such temperatures, baryonic matter could also play an important role in the galactic atmosphere. At a somewhat lower temperature of a few million degrees Kelvin, an atmosphere of 1 GeV protons could generate the $1/r^2$ density profile required for sustaining flat rotation curves. This result is not too dissimilar from early estimates of the temperature of baryonic gas in and around the galactic halo made in the days before the advent of nonbaryonic dark matter.[24],[25] Finally, we remark about a recent observation of a huge reservoir of ionized gas in the circumgalactic medium around the Milky Way radius, extending over a radius of 100 kpc.[26][‡‡] Its mass is reported to be comparable to the total baryonic mass in the disk of the galaxy and its temperature is of the order of 10^6 ^{o}K, which could then possibly sustain the $1/r^2$ density profile necessary for flat rotation curves.

Acknowledgments

We are very grateful to P. Biermann and N. Okada for valuable discussions. A.S. was supported in part by the DOE, Grant No. DE-FG02-10ER41714.

REFERENCES

[1] J. Binney and S. Tremaine "Galactic Dynamics," Princeton, USA: Univ. Pr. (1987) 747 p.

[2] P. J. E. Peebles, "Principles of physical cosmology," Princeton, USA: Univ. Pr. (1993) 718 p.

[3] Emden, R., ' Gaskugeln: Anwendungen der Mechanischen Warmetheorie auf Kosmologische und Meteorologische Probleme', Teubner, Berlin (1907).

[4] S. H. Hansen and R. Piffaretti, "Measuring the dark matter velocity anisotropy in galaxy clusters," arXiv:0705.4680 [astro-ph]; O. Host, S. H. Hansen, R. Piffaretti, A. Morandi, S. Ettori, S. T. Kay and R. Valdarnini, "Measurement of the dark matter velocity anisotropy in galaxy clusters," Astrophys. J. **690**, 358 (2009); S. H. Hansen, A. V. Maccio, E. Romano-Diaz, Y. Hoffman, M. Bruggen, E. Scannapieco and G. S. Stinson, "The temperature of hot gas in galaxies and clusters: baryons dancing to the tune of dark matter," Astrophys. J. **734**, 62 (2011).

[5] J. F. Navarro, C. S. Frenk and S. D. M. White, "A Universal density profile from hierarchical clustering," Astrophys. J. **490**, 493 (1997).

[6] C. R. Keeton, "A catalog of mass models for gravitational lensing," astro-ph/0102341.

[7] L. E. Strigari, J. S. Bullock and M. Kaplinghat, "Determining the Nature of Dark Matter with Astrometry," Astrophys. J. **657**, L1 (2007).

[8] P. Salucci, A. Lapi, C. Tonini, G. Gentile, I. Yegorova and U. Klein, "The Universal Rotation Curve of Spiral Galaxies. 2. The Dark Matter Distribution out to the Virial Radius," Mon. Not. Roy. Astron. Soc. **378**, 41 (2007); F. Donato, G. Gentile, P. Salucci, C. F. Martins, M. I. Wilkinson, G. Gilmore, E. K. Grebel and A. Koch *et al.*, "A constant dark matter halo surface density in galaxies," arXiv:0904.4054 [astro-ph.CO].

[9] C. Destri, H. J. de Vega and N. G. Sanchez, "Fermionic warm dark matter produces galaxy cores in the observed scales," arXiv:1204.3090 [astro-ph.CO].

[10] F. K. Liu, "Polytropic gas spheres: An Approximate analytic solution of the Lane-Emden equation," Mon. Not. Roy. Astron. Soc. **281**, 1197 (1996).

[11] P. Natarajan and D. Lynden-Bell, "An analytic approximation to the isothermal sphere," astro-ph/9604084.

[‡‡] We thank Z. Burrel for pointing out this reference.

[12] Y. Mishchenko and C. -R. Ji, "Molar mass estimate of dark matter from the dark mass distribution measurements," Phys. Rev. D **68**, 063503 (2003).

[13] D. M. Christodoulou and D. Kazanas, "Exact Solutions of the Isothermal Lane-Emden Equation with Rotation and Implications for the Formation of Planets and Satellites," [arXiv:0706.3205 [astro-ph]].

[14] M. Taoso, G. Bertone and A. Masiero, "Dark Matter Candidates: A Ten-Point Test," JCAP **0803**, 022 (2008).

[15] A. Boyarsky, O. Ruchayskiy and D. Iakubovskyi,"A Lower bound on the mass of Dark Matter particles," JCAP **0903**, 005 (2009).

[16] S. Dodelson and L. M. Widrow, "Sterile-neutrinos as dark matter," Phys. Rev. Lett. **72**, 17 (1994).

[17] M. Viel, J. Lesgourgues, M. G. Haehnelt, S. Matarrese and A. Riotto, "Constraining warm dark matter candidates including sterile neutrinos and light gravitinos with WMAP and the Lyman-alpha forest," Phys. Rev. D **71**, 063534 (2005).

[18] P. L. Biermann and A. Kusenko, "Relic keV sterile neutrinos and reionization," Phys. Rev. Lett. **96**, 091301 (2006).

[19] F. Munyaneza and P. L. Biermann, Astron. Astrophys. **458**, L9 (2006).

[20] L. A. Popa and A. Vasile, "Sterile Neutrino as Dark Matter candidate from CMB alone," astro-ph/0701331.

[21] H. J. de Vega and N. G. Sanchez, "Model independent analysis of dark matter points to a particle mass at the keV scale," Mon. Not. Roy. Astron. Soc. **404**, 885 (2010).

[22] C. R. Watson, Z. Li and N. K. Polley, "Constraining Sterile Neutrino Warm Dark Matter with Chandra Observations of the Andromeda Galaxy," JCAP **1203**, 018 (2012).

[23] Y. B. Zeldovich, A. A. Klypin, M. Y. .Khlopov and V. M. Chechetkin, "Astrophysical constraints on the mass of heavy stable neutral leptons," Sov. J. Nucl. Phys. **31**, 664 (1980) [Yad. Fiz. **31**, 1286 (1980)].

[24] Spitzer, L., "On a Possible Interstellar Galactic Corona," ApJ. **124**, 20 (1956).

[25] Kahn, F. D., Woltjer, L., " Intergalactic Matter and the Galaxy," ApJ. **130**, 705 (1959).

[26] A. Gupta, S. Mathur, Y. Krongold, F. Nicastro and M. Galeazzi, "A huge reservoir of ionized gas around the Milky Way: Accounting for the Missing Mass?," arXiv:1205.5037 [astro-ph.HE].